

**Amendments to the Claim:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

**Listing of Claims:**

1. (Previously Presented) Method for prophylaxis or treatment of cancer cachexia in an individual in need of thereof, comprising administration to said individual of a ghrelin-like compound or a pharmaceutically acceptable salt thereof, wherein the ghrelin-like compound comprises a structure defined by formula I

$Z^1 - (X^1)_m - (X^2) - (X^3)_n - Z^2$ , wherein

$Z^1$  is an optionally present protecting group

each  $X^1$  is independently selected from an amino acid, wherein said amino acid is selected from the group consisting of naturally occurring and synthetic amino acids,

$X^2$  is any amino acid selected from the group consisting of naturally occurring and synthetic amino acids, said amino acid being modified with a bulky hydrophobic group

each  $X^3$  is independently selected from an amino acid, wherein said amino acid is selected from the group consisting of naturally occurring and synthetic amino acids,

wherein one or more of  $X^1$  and  $X^3$  optionally may be modified with a bulky hydrophobic group,

$Z^2$  is an optionally present protecting group,

USSN - 10/567,406

m is an integer in the range of from 1-10

n is 0 or an integer in the range of from 1-35,

and wherein:

(a) said ghrelin-like compound or pharmaceutically acceptable salt thereof is 27-28 amino acids in length, with the proviso that said ghrelin-like compound is at least 80 % homologous to SEQ ID NO 1, such as at least 85 % homologous to SEQ ID NO: 1

and/or

(b) said ghrelin-like compound is at least 90 % homologous to SEQ ID NO 1.

2. (Previously Presented) The method according to claim 1, wherein said ghrelin-like compound is at least 95 % homologous to SEQ ID NO 1.

3. (Previously Presented) The method according to claim 1, wherein said ghrelin-like compound is at least 98 % homologous to SEQ ID NO 1.

4. (Cancelled).

5. (Previously Presented) The method according to claim 1, wherein  $X^2$  is selected from the group consisting of modified Ser, modified Cys and modified Lys.

6. (Previously Presented) The method according to claim 1, wherein the ghrelin-like compound is selected from a compound of

formula II  $Z^1 - \text{Gly} - (X^1)_{m-1} - (X^2) - (X^3)_n - Z^2$ ,

formula III  $Z^1 - \text{Gly} - \text{Ser} - (X^2) - (X^3)_n - Z^2$ , and

formula IV  $Z^1 - \text{Gly} - (X^2) - (X^3)_n - Z^2$ .

7-12. (Cancelled).

13. (Previously Presented) The method according to claim 1, wherein the ghrelin-like compound is ghrelin or a pharmaceutically acceptable salt thereof.

14. (Currently Amended) The method according to ~~to~~ claim 1, wherein the ghrelin-like compound has SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3.

15. (Previously Presented) The method according to claim 1, wherein the medicament is in a formulation for subcutaneous administration.

16. (Previously Presented) The method according to claim 1, wherein the formulation comprises the ghrelin-like compound or a salt thereof as a lyophilisate and the formulation further comprises a solvent, said lyophilisate and said solvent being in separate compartments until administration.

17. (Previously Presented) The method according to claim 1, wherein the formulation is a solution of the ghrelin-like compound or a salt thereof.

18. (Previously Presented) The method according to claim 16 wherein the solvent is saline.

19. (Previously Presented) The method according to claim 1, wherein the medicament is administered prior to or during a meal.

20. (Previously Presented) The method according to claim 1, wherein the medicament is administered in a concentration equivalent to from 10 ng to 10 mg ghrelin per kg bodyweight.

21. (Cancelled).

22. (Previously Presented) The method according to claim 1, wherein the medicament is administered in a concentration equivalent to from 0.1  $\mu$ g to 1 mg ghrelin per kg bodyweight.

23. (Previously Presented) The method according to claim 1, wherein the medicament is administered as a bolus prior to or during a meal, said bolus comprising an amount of the

ghrelin-like compound or a salt thereof equivalent to from 0.3  $\mu$ g to 600 mg ghrelin.

24-26. (Cancelled).

27. (Previously Presented) The method according to claim 1, wherein the cancer cachexia is caused by a catabolic disorder.

28. (Previously Presented) The method according to claim 1, wherein the cancer cachexia is caused by an anorectic disorder.

29. (Previously Presented) The method according to claim 1, where the individual is suffering from a cancer selected from the group consisting of lung cancer, pancreatic cancer, liver cancer, and GI tract cancers.

30. (Cancelled).

31. (Previously Presented) The method according to claim 1, wherein the treatment or prevention of cancer cachexia leads to stimulation of appetite, stimulation of food intake, stimulation of weight gain or weight maintenance, and/or increased body fat mass.

32. (Cancelled).

33. (Withdrawn - Currently Amended) ~~A The method of claim 1, for preventing or treating cancer, further comprising administering to an subjecting the individual in need thereof an effective amount of a ghrelin-like compound in combination with to an anti-neoplastic treatment.~~

34. (Withdrawn) The method according to claim 33, wherein the antineoplastic treatment is radiotherapy.

35. (Withdrawn) The method according to claim 33, wherein the antineoplastic treatment is chemotherapy.

36. (Cancelled).

37. (Currently Amended) ~~A The method of claim 1, for preventing or treating cachexia, further comprising administering to an individual in need thereof administration of an effective amount of an a ghrelin-like compound and an effective amount of ~~a NSAID medicament.~~~~

38. (Cancelled).

39. (Currently Amended) A method for stimulation of appetite in an individual, comprising administering to the individual a subcutaneous dosage of a medicament comprising a ghrelin-like compound or a pharmaceutically acceptable salt thereof, in a formulation for subcutaneous administration,

wherein the ghrelin-like compound comprises a structure defined by formula I

$Z^1 - (X^1)_m - (X^2) - (X^3)_n - Z^2$ , wherein

$Z^1$  is an optionally present protecting group

each  $X^1$  is independently selected from an amino acid, wherein said amino acid is selected from the group consisting of naturally occurring and synthetic amino acids,

$X^2$  is any amino acid selected from the group consisting of naturally occurring and synthetic amino acids, said amino acid being modified with a bulky hydrophobic group,

each  $X^3$  is independently selected from an amino acid, wherein said amino acid is selected from the group consisting of naturally occurring and synthetic amino acids,

wherein one or more of  $X^1$  and  $X^3$  optionally may be modified with a bulky hydrophobic group, preferably an acyl group, or a fatty acid,

$Z^2$  is an optionally present protecting group,

USSN - 10/567,406

m is an integer in the range of from 1-10

n is 0 or an integer in the range of from 1-35

and wherein

(a) said ghrelin-like compound or pharmaceutically acceptable salt thereof is 27-28 amino acids in length, with the proviso that said ghrelin-like compound is at least 80 % homologous to SEQ ID NO 1, such as at least 85 % homologous to SEQ ID NO: 1

and/or

(b) said ghrelin-like compound is at least 90 % homologous to SEQ ID NO 1.

40. (Previously Presented) The method according to claim 39, wherein the ghrelin-like compound has SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3.

41-42. (Cancelled).

43. (Previously Presented) The method according to claim 39, wherein the medicament is administered in a concentration equivalent to from 10 ng to 10 mg ghrelin per kg bodyweight.

44. (Previously Presented) The method according to claim 39, wherein said individual is suffering from lipodystrophy.